



NSAI
Standards

Irish Standard Recommendation
S.R. CEN/TR 16998:2016

Ambient air - Report on nitro- and oxy-PAHs - Origin, toxicity, concentrations and measurement methods

S.R. CEN/TR 16998:2016

Incorporating amendments/corrigenda/National Annexes issued since publication:

The National Standards Authority of Ireland (NSAI) produces the following categories of formal documents:

I.S. xxx: Irish Standard — national specification based on the consensus of an expert panel and subject to public consultation.

S.R. xxx: Standard Recommendation — recommendation based on the consensus of an expert panel and subject to public consultation.

SWiFT xxx: A rapidly developed recommendatory document based on the consensus of the participants of an NSAI workshop.

This document replaces/revises/consolidates the NSAI adoption of the document(s) indicated on the CEN/CENELEC cover/Foreword and the following National document(s):

NOTE: The date of any NSAI previous adoption may not match the date of its original CEN/CENELEC document.

This document is based on:

CEN/TR 16998:2016

Published:

2016-11-30

This document was published under the authority of the NSAI and comes into effect on:

2016-12-18

ICS number:

13.040.20

NOTE: If blank see CEN/CENELEC cover page

NSAI
1 Swift Square,
Northwood, Santry
Dublin 9

T +353 1 807 3800
F +353 1 807 3838
E standards@nsai.ie
W NSAI.ie

Sales:
T +353 1 857 6730
F +353 1 857 6729
W standards.ie

Údarás um Chaighdeáin Náisiúnta na hÉireann

National Foreword

S.R. CEN/TR 16998:2016 is the adopted Irish version of the European Document CEN/TR 16998:2016, Ambient air - Report on nitro- and oxy-PAHs - Origin, toxicity, concentrations and measurement methods

This document does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

Compliance with this document does not of itself confer immunity from legal obligations.

In line with international standards practice the decimal point is shown as a comma (,) throughout this document.

This page is intentionally left blank

TECHNICAL REPORT

CEN/TR 16998

RAPPORT TECHNIQUE

TECHNISCHER BERICHT

November 2016

ICS 13.040.20

English Version

Ambient air - Report on nitro- and oxy-PAHs - Origin, toxicity, concentrations and measurement methods

Air ambiant - Rapport sur les nitro- et oxy-HAP -
Origine, toxicité, concentrations et méthodes de
mesure

Außenluft - Bericht über Nitro- und Oxy-PAHs -
Herkunft, Toxizität, Konzentrationen und
Messverfahren

This Technical Report was approved by CEN on 21 October 2016. It has been drawn up by the Technical Committee CEN/TC 264.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

CEN/TR 16998:2016 (E)

Contents		Page
European foreword		3
Introduction		4
1	Scope	5
2	Symbols and abbreviations	5
3	Literature overview	5
4	Conclusions	20
5	Recommendations	20
Annex A (informative) Sampling and analysis by GC-MS of some nitro- and oxy-PAHs associated to ambient particulate matter		22
A.1	Sampling	22
A.2	Analytical materials	22
A.2.1	Glassware and sample handling	22
A.2.2	Reagents and Solvents	22
A.2.3	Extraction apparatus and materials	23
A.2.4	Evaporation apparatus and materials	24
A.2.5	Clean-up Material	24
A.2.6	Weighting Apparatus	24
A.2.7	Analytical system	24
A.3	Extraction	24
A.4	Clean-up	24
A.5	Analysis	25
A.6	Results	25
A.7	Quality assurance	30
Annex B (informative) Carcinogenicity and references to nitro- and oxy-PAHs		31
Annex C (informative) Mutagenicity of nitro-PAHs		34
Annex D (informative) Diesel exhaust data		35
Annex E (informative) Structures of nitro- and oxy-PAHs referred in this Technical Report		36
Bibliography		43

European foreword

This document (CEN/TR 16998:2016) has been prepared by Technical Committee CEN/TC 264 “Air quality”, the secretariat of which is held by DIN.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

CEN/TR 16998:2016 (E)

Introduction

Nitro-PAHs and oxy-PAHs are found in ambient air samples and there are strong indications that they are as harmful as PAHs. Several compounds are classified as probably carcinogenic for humans (see Table in Annex A) and nitro-PAHs are reported to be strongly mutagenic. Photooxidation of volatile PAHs gives rise to the formation of secondary aerosols (Chan et al. 2009, Kautzman et al. 2010, Shakya and Griffin, 2010).

1-Nitropyrene and 2-nitrofluorene are discussed as marker compounds for diesel exhaust and other combustion processes. 2-Nitropyrene and 2-nitrofluoranthene are good marker substances for the formation of nitro-PAHs by secondary reactions.

This Technical Report presents the state of the art of the oxy- and nitro-PAHs topics.

1 Scope

This Technical Report is focused on the presence of nitro- and oxy-PAHs in ambient air. It describes how nitro- and oxy-PAH are formed, what typical concentrations are found, what is known about their toxicity, and what sampling and measurement techniques are available.

The conclusions of this report are that nitro- and oxy-PAHs concentrations are present in the atmosphere in levels that are of concern regarding their high toxicity. Information on the presence of these compounds in ambient air is as relevant as information about PAHs. Validated techniques for the measurement of nitro- and oxy-PAHs are available.

2 Symbols and abbreviations

DNA	Deoxyribonucleic acid
EI	Electron ionization
CD	Chemiluminescence detection
FD	Fluorescence detection
GC-MS	Gas chromatography – mass spectrometry
GC-NICI-MS	Gas chromatography – negative ion chemical ionization – mass spectrometry
HPLC	High performance liquid chromatography
HPLC-FD	HPLC – fluorescence detection
HPLC-CD	HPLC – chemiluminescence detection
IARC	International Agency for Research on Cancer
LC	Liquid chromatography
MS	Mass spectrometry
NICI	Negative ion chemical ionization
Nitro-PAHs	Nitrated polycyclic aromatic hydrocarbons
Oxy-PAHs	Oxygenated polycyclic aromatic hydrocarbons
PAHs	Polycyclic aromatic hydrocarbons
SPE	Solid phase extraction
ToF-MS	Time of flight mass spectrometry

3 Literature overview

3.1 Nitro-PAHs

3.1.1 Sources

3.1.1.1 General

Nitro-PAHs in the atmosphere originate mainly from combustion sources and are produced from both gas and heterogeneous phase reactions of the parent PAHs with atmospheric oxidants such as NO₃, N₂O₅, O₃, OH and peroxide radicals (Arey et al., 1986; Atkinson et al., 1990; Keyte et al., 2013; Pitts et al., 1985; Pitts Jr et al., 1978) in the presence of nitrogen oxides.

This is a free preview. Purchase the entire publication at the link below:

[Product Page](#)

-
- [Looking for additional Standards? Visit Intertek Inform Infostore](#)
 - [Learn about LexConnect, All Jurisdictions, Standards referenced in Australian legislation](#)
-