



NSAI
Standards

Irish Standard
I.S. EN ISO 18465:2017

Microbiology of the food chain - Quantitative determination of emetic toxin (cereulide) using LC-MS/MS (ISO 18465:2017)

I.S. EN ISO 18465:2017

Incorporating amendments/corrigenda/National Annexes issued since publication:

The National Standards Authority of Ireland (NSAI) produces the following categories of formal documents:

I.S. xxx: Irish Standard — national specification based on the consensus of an expert panel and subject to public consultation.

S.R. xxx: Standard Recommendation — recommendation based on the consensus of an expert panel and subject to public consultation.

SWiFT xxx: A rapidly developed recommendatory document based on the consensus of the participants of an NSAI workshop.

This document replaces/revises/consolidates the NSAI adoption of the document(s) indicated on the CEN/CENELEC cover/Foreword and the following National document(s):

NOTE: The date of any NSAI previous adoption may not match the date of its original CEN/CENELEC document.

This document is based on:

EN ISO 18465:2017

Published:

2017-02-08

*This document was published
under the authority of the NSAI
and comes into effect on:*

2017-02-27

ICS number:

07.100.30

NOTE: If blank see CEN/CENELEC cover page

NSAI
1 Swift Square,
Northwood, Santry
Dublin 9

T +353 1 807 3800
F +353 1 807 3838
E standards@nsai.ie
W NSAI.ie

Sales:
T +353 1 857 6730
F +353 1 857 6729
W standards.ie

Údarás um Chaighdeáin Náisiúnta na hÉireann

National Foreword

I.S. EN ISO 18465:2017 is the adopted Irish version of the European Document EN ISO 18465:2017, Microbiology of the food chain - Quantitative determination of emetic toxin (cereulide) using LC-MS/MS (ISO 18465:2017)

This document does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

Compliance with this document does not of itself confer immunity from legal obligations.

In line with international standards practice the decimal point is shown as a comma (,) throughout this document.

This page is intentionally left blank

EUROPEAN STANDARD
NORME EUROPÉENNE
EUROPÄISCHE NORM

EN ISO 18465

February 2017

ICS 07.100.30

English Version

**Microbiology of the food chain - Quantitative
determination of emetic toxin (cereulide) using LC-MS/MS
(ISO 18465:2017)**

Microbiologie de la chaîne alimentaire - Détermination
quantitative de la toxine émétique (céréulide) par CL-
SM/SM (ISO 18465:2016)

Mikrobiologie der Lebensmittelkette - Quantitative
Bestimmung von emetischem Toxin (Cereulid) mittels
LC-MS/MS (ISO 18465:2017)

This European Standard was approved by CEN on 28 December 2016.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

Contents	Page
European Foreword.....	3

European Foreword

This document (EN ISO 18465:2017) has been prepared by Technical Committee CEN/TC 275 "Food analysis - Horizontal methods", the secretariat of which is held by DIN, in collaboration with Technical Committee ISO/TC 34 "Food products".

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by August 2017 and conflicting national standards shall be withdrawn at the latest by August 2017.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

This document has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

Endorsement notice

The text of ISO 18465:2017 has been approved by CEN as EN ISO 18465:2017 without any modification.

This page is intentionally left blank

INTERNATIONAL STANDARD

**ISO
18465**

First edition
2017-01

Microbiology of the food chain — Quantitative determination of emetic toxin (cereulide) using LC-MS/MS

*Microbiologie de la chaîne alimentaire — Détermination quantitative
de la toxine émétique (céréulide) par CL-SM/SM*



Reference number
ISO 18465:2017(E)

© ISO 2017

ISO 18465:2017(E)



COPYRIGHT PROTECTED DOCUMENT

© ISO 2017, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Ch. de Blandonnet 8 • CP 401
CH-1214 Vernier, Geneva, Switzerland
Tel. +41 22 749 01 11
Fax +41 22 749 09 47
copyright@iso.org
www.iso.org

Contents

Page

Foreword	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Terms and definitions	1
4 General principle	1
5 Reagents	1
6 Apparatus and equipment	3
7 Procedure	4
7.1 Sample preparation	4
7.2 Standard preparation	4
7.3 LC-MS analysis.....	5
7.3.1 LC conditions.....	5
7.3.2 MS conditions and tuning parameters.....	5
7.3.3 Transitions (multiple reaction monitoring, MRM).....	6
8 Calculation	6
9 Quality controls	7
10 Precision	8
10.1 General.....	8
10.2 Repeatability.....	8
10.3 Reproducibility.....	9
Annex A (informative) Results of the interlaboratory study	10
Annex B (informative) Possible transitions of cereulide in MS^[3]	12
Bibliography	14

ISO 18465:2017(E)

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: www.iso.org/iso/foreword.html.

This document was prepared by the European Committee for Standardization (CEN) Technical Committee CEN/TC 275, *Food Analysis — Horizontal methods*, in collaboration with ISO Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 9, *Microbiology*, in accordance with the agreement on technical cooperation between ISO and CEN (Vienna Agreement).

Introduction

Cereulide, the emetic toxin produced in foods by certain strains of *Bacillus cereus*, is a heat and acid stable toxin that causes nausea and vomiting when ingested. In very rare cases, people can die after ingestion of the toxin. Due to its stability, the toxin may still be present even when *B. cereus* can no longer be detected. The presence of cereulide seems to be linked to starch-rich foods like rice (dishes) and pasta (dishes). However, recent data suggest that the occurrence of food borne outbreaks due to cereulide is more common to foods in general[9]. The toxin has a cyclic structure and consists of in total 12 monomers as a repeat of (D-O-Leucine-D-Alanine-L-O-Valine-L-Valine). Several methods have been developed for the detection and/or quantification of the toxin. Some of these methods are nonspecific bio-assays[3, 4] and other methods are specifically based on the chemical analysis using liquid chromatography with mass spectrometry (LC-MS/MS) for the detection and quantification of the toxin[5, 6, 7, 8]. The chemical methods are more specific for cereulide and have, therefore, been chosen as the starting point for standardization of a method for the quantification of cereulide. Recently, research has been done for the chemodiversity of cereulide. At least 18 cereulide variants were detected by UHPLC-TOFMS and ion-trap MSⁿ sequencing, among which the previously unknown isocereulides A-G[10].

Microbiology of the food chain — Quantitative determination of emetic toxin (cereulide) using LC-MS/MS

1 Scope

This document describes the quantitative analysis of the emetic toxin cereulide using high performance liquid chromatography (HPLC) or ultra performance liquid chromatography (UHPLC) connected to a tandem mass spectrometer (LC-MS/MS).

This document is applicable to the analysis of the toxin in products intended for human consumption.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 3696, *Water for analytical laboratory use — Specification and test methods*

ISO 1042, *Laboratory glassware — One-mark volumetric flasks*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>
- ISO Online browsing platform: available at <http://www.iso.org/obp>

3.1

cereulide

toxin cyclo[D-O-Leucine-D-Alanine-L-O-Valine-L-Valine]₃ produced by certain strains of the species of *B. cereus*

4 General principle

Cereulide is extracted from the food matrix by shaking the sample with acetonitrile. ¹³C₆-Cereulide is used as an internal standard. The components in the solution are separated using HPLC or UHPLC and subsequently detected using tandem mass spectrometry (LC-MS/MS). For MS, the electro spray ionization technique (ESI) is used, using the positive mode. The level of emetic toxin (cereulide) is expressed as µg cereulide/kg product.

5 Reagents

Use only reagents of recognized analytical grade, unless otherwise specified.

5.1 Water, according to ISO 3696.

5.2 Acetonitrile, LC-MS grade.

This is a free preview. Purchase the entire publication at the link below:

[Product Page](#)

-
- Looking for additional Standards? Visit Intertek Inform Infostore
 - Learn about LexConnect, All Jurisdictions, Standards referenced in Australian legislation
-