



NSAI
Standards

Irish Standard
I.S. EN 14103:2020

Fat and oil derivatives - Fatty Acid Methyl Esters (FAME) - Determination of ester and linolenic acid methyl ester contents

I.S. EN 14103:2020

Incorporating amendments/corrigenda/National Annexes issued since publication:

The National Standards Authority of Ireland (NSAI) produces the following categories of formal documents:

I.S. xxx: Irish Standard — national specification based on the consensus of an expert panel and subject to public consultation.

S.R. xxx: Standard Recommendation — recommendation based on the consensus of an expert panel and subject to public consultation.

SWiFT xxx: A rapidly developed recommendatory document based on the consensus of the participants of an NSAI workshop.

This document replaces/revises/consolidates the NSAI adoption of the document(s) indicated on the CEN/CENELEC cover/Foreword and the following National document(s):

NOTE: The date of any NSAI previous adoption may not match the date of its original CEN/CENELEC document.

This document is based on:

EN 14103:2020

Published:

2020-01-22

This document was published under the authority of the NSAI and comes into effect on:

2020-02-09

ICS number:

67.200.10

NOTE: If blank see CEN/CENELEC cover page

NSAI
1 Swift Square,
Northwood, Santry
Dublin 9

T +353 1 807 3800
F +353 1 807 3838
E standards@nsai.ie
W NSAI.ie

Sales:
T +353 1 857 6730
F +353 1 857 6729
W standards.ie

Údarás um Chaighdeáin Náisiúnta na hÉireann

National Foreword

I.S. EN 14103:2020 is the adopted Irish version of the European Document EN 14103:2020, Fat and oil derivatives - Fatty Acid Methyl Esters (FAME) - Determination of ester and linolenic acid methyl ester contents

This document does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

For relationships with other publications refer to the NSAI web store.

Compliance with this document does not of itself confer immunity from legal obligations.

In line with international standards practice the decimal point is shown as a comma (,) throughout this document.

This page is intentionally left blank

CEN/TC 307

Date: 2018-05

prEN 14103:2018

Secretariat: AFNOR

Fat and oil derivatives — Fatty Acid Methyl Esters (FAME) — Determination of ester and linolenic acid methyl ester contents

Erzeugnisse aus pflanzlichen und tierischen Fetten und Ölen — Fettsäure-Methylester (FAME) — Bestimmung des Ester-Gehaltes und des Gehaltes an Linolensäure-Methylester

Produits dérivés des corps gras — Esters méthyliques d'acides gras — Détermination de la teneur en ester et en ester méthylique de l'acide linoléique

ICS:

3

Contents	Page
European foreword	3
1 Scope.....	4
2 Normative references.....	4
3 Terms and definitions	4
4 Principle	4
5 Reagents	5
6 Apparatus	5
7 Sampling.....	5
8 Procedure	5
8.1 Operating conditions	5
8.2 Internal standard GC purity determination	6
8.3 FAME sample preparation and analysis	6
8.4 Identification	7
9 Calculation.....	7
9.1 Determination of ester content	7
9.2 Determination of linolenic acid methyl ester.....	10
9.3 Expression of results	10
10 Precision	10
10.1 General.....	10
10.2 Repeatability	10
10.3 Reproducibility.....	11
11 Test report.....	11
Annex A (informative) Chromatograms of FAME samples - Determination of ester content and linolenic acid methyl ester content – Summary of the analytical conditions.....	12
Annex B (informative) Results of an interlaboratory trial	16
Annex C (informative) Calculation of the pattern of fatty acid methyl esters.....	18
Bibliography	19

European foreword

This document (EN 14103:2020) has been prepared by Technical Committee CEN/TC 307 “Oilseeds, vegetable and animal fats and oils and their by-products — Methods of sampling and analysis”, the secretariat of which is held by AFNOR.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by July 2020, and conflicting national standards shall be withdrawn at the latest by July 2020.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN shall not be held responsible for identifying any or all such patent rights.

This document supersedes EN 14103:2011.

In comparison with the previous edition, the following technical modifications have been made:

- a) note on natural nonadecanoic acid methyl ester added in the scope;
- b) new procedure to check nonadecanoic acid methyl ester purity, with new GC conditions, and reduction of the minimum GC purity (99,5 to 99,0 % (*m/m*));
- c) calculation of results revised by incorporation of theoretical flame ionization detector correction factor (TCF), this gives a better accuracy of the calculated contents in case of presence of methyl esters with short chains;
- d) new inter-laboratories study (ILS) conducted and precision adopted;
- e) new sample chromatograms recorded and added;
- f) calculation of the pattern of fatty acid methyl esters incorporated as informative Annex C;
- g) modification of the way of integration in 8.4 by taking all the peaks into consideration whereas in the previous edition all the peaks identified as fatty acid methyl esters were taken into consideration;
- h) increase of the FAME sample test portion in 8.3 to 250 mg whereas in the previous edition the sample test portion was 100 mg;
- i) document revised editorially.

According to the CEN-CENELEC Internal Regulations, the national standards organisations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Republic of North Macedonia, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

EN 14103:2020 (E)

1 Scope

The purpose of this document is to describe a procedure for the determination of the ester content in fatty acid methyl esters (FAME) intended for incorporation into diesel oil. It also allows determining the linolenic acid methyl ester content. It allows verifying that the ester content of FAME is greater than 90 % (*m/m*) and that the linolenic acid methyl ester content is between 1 % (*m/m*) and 15 % (*m/m*).

The precision was established using FAMES with an ester content of 95 % (*m/m*) and 100 % (*m/m*) only, thus covering the range of the limit value. The method is also suitable outside of this range; however, precision for lower concentrations is subject to further work.

This method is suitable for FAME which contains methyl esters between C6 and C24.

NOTE 1 For the purposes of this document, the term “% (*m/m*)” is used to represent the mass fractions.

This method was elaborated for FAME samples from usual raw material. For FAME sample from unidentified raw material, a solution of the test sample is prepared without any internal standard addition, in order to verify the absence of natural nonadecanoic acid methyl ester or other unknown substances co-eluting with the IS.

NOTE 2 The calculation method of the pattern of fatty acid methyl esters is given in Annex C.

WARNING — The use of this method may involve hazardous equipment, materials and operations. This method does not purport to address to all of the safety problems associated with its use, but it is the responsibility of the user to search and establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use.

2 Normative references

There are no normative references in this document.

3 Terms and definitions

No terms and definitions are listed in this document.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>
- ISO Online browsing platform: available at <http://www.iso.org/obp>

4 Principle

Determination of the mass percentage of total methyl esters of fatty acids and the mass percentage of linolenic acid methyl ester present in the sample, by gas chromatography according to a procedure using internal calibration (nonadecanoic acid methyl ester).

This is a free preview. Purchase the entire publication at the link below:

[Product Page](#)

-
- [Looking for additional Standards? Visit Intertek Inform Infostore](#)
 - [Learn about LexConnect, All Jurisdictions, Standards referenced in Australian legislation](#)
-