



National Standards Authority of Ireland

STANDARD RECOMMENDATION

S.R. CEN/TR 15547:2007

ICS 13.040.30

**WORKPLACE ATMOSPHERES -  
CALCULATION OF THE HEALTH-RELATED  
AEROSOL FRACTION CONCENTRATION  
FROM THE CONCENTRATION MEASURED BY  
A SAMPLER WITH KNOWN PERFORMANCE  
CHARACTERISTICS**

National Standards  
Authority of Ireland  
Glasnevin, Dublin 9  
Ireland

Tel: +353 1 807 3800  
Fax: +353 1 807 3838  
<http://www.nsai.ie>

**Sales**  
<http://www.standards.ie>

*This Standard  
Recommendation was  
published under the authority  
of the National Standards  
Authority of Ireland and  
comes into effect on:  
16 March 2007*

**NO COPYING WITHOUT NSAI  
PERMISSION EXCEPT AS  
PERMITTED BY COPYRIGHT  
LAW**

© NSAI 2007

**Price Code F**

Údarás um Chaighdeáin Náisiúnta na hÉireann



TECHNICAL REPORT  
RAPPORT TECHNIQUE  
TECHNISCHER BERICHT

**CEN/TR 15547**

February 2007

---

ICS 13.040.30

English Version

**Workplace atmospheres - Calculation of the health-related  
aerosol fraction concentration from the concentration measured  
by a sampler with known performance characteristics**

Atmosphères des lieux de travail - Calcul de la  
concentration en fractions d'aérosols liées à la santé à  
partir de la concentration mesurée à l'aide d'un dispositif de  
prélèvement ayant des caractéristiques de performances  
connues

Arbeitsplatzatmosphäre - Berechnung der  
gesundheitsbezogenen Fraktion der Aerosolkonzentration  
anhand der mit einem Probenahmegerät mit bekannten  
Leistungseigenschaften gemessenen Konzentration

This Technical Report was approved by CEN on 4 December 2006. It has been drawn up by the Technical Committee CEN/TC 137.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION  
COMITÉ EUROPÉEN DE NORMALISATION  
EUROPÄISCHES KOMITEE FÜR NORMUNG

**Management Centre: rue de Stassart, 36 B-1050 Brussels**

<b>Contents</b>		Page
<b>Foreword</b> .....		<b>3</b>
<b>Introduction</b> .....		<b>4</b>
<b>1</b>	<b>Scope</b> .....	<b>5</b>
<b>2</b>	<b>Normative references</b> .....	<b>5</b>
<b>3</b>	<b>Method to produce improved concentration data</b> .....	<b>5</b>
<b>4</b>	<b>Data needed to calculate the concentration of a health-related aerosol fraction</b> .....	<b>6</b>
<b>5</b>	<b>Calculation method</b> .....	<b>6</b>
<b>Annex A (informative) Numerical example of calculation</b> .....		<b>9</b>
<b>Annex B (informative) Field sampling example</b> .....		<b>12</b>
<b>Bibliography</b> .....		<b>14</b>

## **Foreword**

This document (CEN/TR 15547:2007) has been prepared by Technical Committee CEN/TC 137 “Assessment of workplace exposure to chemical and biological agents”, the secretariat of which is held by DIN.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN shall not be held responsible for identifying any or all such patent rights.

## **CEN/TR 15547:2007 (E)**

### **Introduction**

Exposure assessment of workers to particulate matter dispersed into the air at the workplace is generally achieved through aerosol sampling by using instruments designed for measuring health-related aerosol fractions as defined in EN 481. EN 13205 gives a methodology to evaluate sampler performance. The knowledge of the sampling efficiency of a sampler is used to calculate the bias and the accuracy in concentration for log-normally distributed aerosols. Bias and accuracy maps give an overall indication on sampler performance when sampling health-related aerosol fractions. This performance varies with particle size distribution of sampled aerosol.

Many different samplers can be used for the same purpose, depending on local circumstances or the current practice in the country where these measurements have to be performed. Even with samplers whose performances are quite similar, some significant differences in measured concentrations can occur between these samplers, depending on the aerosol measured. Furthermore, the concentration measured by a sampler is not actually the conventional concentration even if the sampler fulfils the performance criteria stated in EN 13205. This is due to the fact that the particle-size selectivity of the sampler does not generally coincide exactly with the conventional sampling curve over the whole particle-size range.

In the revision of EN 482 presently under way, the uncertainty estimate of a measurement procedure should be expanded to meet the requirements of ENV 13005 complying with GUM (ISO Guide to the expression of Uncertainty in Measurements). This requires that all uncertainties encountered by the use of a measurement procedure (except interlaboratory variation) have to be accounted for. For the special case of aerosol sampling this means that the uncertainty in an expected bias of the sample due to non-ideal collection characteristics will only be estimated for a very wide range of size distributions. The calculations presented in this Technical Report can help to significantly reduce this uncertainty by first estimating a restricted range of size distributions in which the sampler was actually used, and then estimate the bias uncertainty only over this narrow range of size distributions. For an aerosol sampler, the variability due to bias is in many cases a major component of the uncertainty.

This is a free preview. Purchase the entire publication at the link below:

[Product Page](#)

- 
- [Looking for additional Standards? Visit Intertek Inform Infostore](#)
  - [Learn about LexConnect, All Jurisdictions, Standards referenced in Australian legislation](#)
-