



NSAI
Standards

Irish Standard
I.S. EN ISO 16198:2015

Soil quality - Plant-based test to assess the environmental bioavailability of trace elements to plants (ISO 16198:2015)

I.S. EN ISO 16198:2015

Incorporating amendments/corrigenda/National Annexes issued since publication:

The National Standards Authority of Ireland (NSAI) produces the following categories of formal documents:

I.S. xxx: Irish Standard — national specification based on the consensus of an expert panel and subject to public consultation.

S.R. xxx: Standard Recommendation — recommendation based on the consensus of an expert panel and subject to public consultation.

SWiFT xxx: A rapidly developed recommendatory document based on the consensus of the participants of an NSAI workshop.

This document replaces/revises/consolidates the NSAI adoption of the document(s) indicated on the CEN/CENELEC cover/Foreword and the following National document(s):

NOTE: The date of any NSAI previous adoption may not match the date of its original CEN/CENELEC document.

This document is based on:

EN ISO 16198:2015

Published:

2015-01-14

This document was published under the authority of the NSAI and comes into effect on:

2015-02-24

ICS number:

13.080.30

NOTE: If blank see CEN/CENELEC cover page

NSAI
1 Swift Square,
Northwood, Santry
Dublin 9

T +353 1 807 3800
F +353 1 807 3838
E standards@nsai.ie
W NSAI.ie

Sales:
T +353 1 857 6730
F +353 1 857 6729
W standards.ie

Údarás um Chaighdeáin Náisiúnta na hÉireann

EUROPEAN STANDARD

EN ISO 16198

NORME EUROPÉENNE

EUROPÄISCHE NORM

January 2015

ICS 13.080.30

English Version

Soil quality - Plant-based test to assess the environmental bioavailability of trace elements to plants (ISO 16198:2015)

Qualité du sol - Test végétal pour l'évaluation de la biodisponibilité environnementale des éléments traces pour les végétaux (ISO 16198:2015)

Bodenbeschaffenheit - Pflanzenbasierter Test zur Beurteilung der umweltrelevanten Bioverfügbarkeit von Spurenelementen für Pflanzen (ISO 16198:2015)

This European Standard was approved by CEN on 24 October 2014.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

EN ISO 16198:2015 (E)

Contents	Page
Foreword.....	3

Foreword

This document (EN ISO 16198:2015) has been prepared by Technical Committee ISO/TC 190 "Soil quality" in collaboration with Technical Committee CEN/TC 345 "Characterization of soils" the secretariat of which is held by NEN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by July 2015, and conflicting national standards shall be withdrawn at the latest by July 2015.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

Endorsement notice

The text of ISO 16198:2015 has been approved by CEN as EN ISO 16198:2015 without any modification.

This page is intentionally left blank

INTERNATIONAL STANDARD

ISO
16198

First edition
2015-01-15

Soil quality — Plant-based test to assess the environmental bioavailability of trace elements to plants

*Qualité du sol — Test végétal pour l'évaluation de la biodisponibilité
environnementale des éléments traces pour les végétaux*



Reference number
ISO 16198:2015(E)

© ISO 2015

ISO 16198:2015(E)



COPYRIGHT PROTECTED DOCUMENT

© ISO 2015

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Contents

Page

Foreword	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Terms and definitions	2
4 Principle	3
5 Laboratory apparatus	5
6 Reagents	5
6.1 General.....	5
7 Biological and growing apparatus	6
7.1 Plant species.....	6
7.2 Biotest apparatus.....	6
7.3 Composition of the nutrient solutions.....	9
7.4 Climatic conditions in the growth chamber.....	10
8 Pre-treatment and analysis of soil or soil material sample	10
8.1 Sample size and particle size reduction.....	10
8.2 Analyses.....	10
9 Experimental and analytical procedure	11
9.1 Overview of the procedure.....	11
9.2 Selection and preparation of seeds.....	11
9.3 Preculture period: Germination and pre-growth in hydroponics.....	11
9.4 Preparation and incubation of soil or soil material.....	12
9.5 Test culture period: Plant growth in contact with soil or soil material.....	12
9.6 Plant harvests.....	13
9.7 Grinding and digestion of shoots and roots.....	13
9.8 Analytical determination.....	14
10 Validity criteria	14
11 Assessment of the results	14
11.1 Determination of trace element concentrations and uptake flux in plants.....	14
11.2 Data presentation.....	16
11.3 Expression of the results.....	16
12 Statistical analysis	16
12.1 General.....	16
12.2 Plant biomasses.....	16
12.3 Bioavailability end points.....	17
13 Test report	17
Annex A (informative) Plant species adapted to the biotest procedure	19
Annex B (informative) Technical drawings of the different components of the biotest	21
Annex C (informative) Seed selection and seed density in plant pot for a range of species tested with the standardized experimental procedure	24
Annex D (informative) Digestion and analysis of plant samples	26
Annex E (informative) Range of biomasses and trace element quantities in control plant pots	28
Annex F (informative) Ring-test	29
Bibliography	43

ISO 16198:2015(E)

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the WTO principles in the Technical Barriers to Trade (TBT) see the following URL: Foreword - Supplementary information

The committee responsible for this document is ISO/TC 190, *Soil quality*, Subcommittee SC 7, *Soil and site assessment*.

Introduction

One of the main objectives of ISO 17402 is to define a conceptual framework of the bioavailability of contaminants in soils and soil materials, and to provide a guidance for the selection of methods able to be standardized for the measurement of bioavailability. Bioavailability was thus defined according to three successive steps:

- a) “environmental availability”;
- b) “environmental bioavailability”;
- c) “toxicological bioavailability”.

The environmental bioavailability is consequently a prerequisite to the assessment of the toxicological bioavailability and is directly related to the impact of pollutants on major functions of soil in the ecosystem and more particularly to habitat and retention functions.

Environmental bioavailability can be estimated with either chemical or biological methods. In the case of trace elements, chemical methods are usually the cheapest, easy to perform, and some of them are already standardized at national or international level (e.g. ISO 19730). However, chemical methods which, strictly speaking, measure the environmental availability in soils have to be correlated with biological measurements before being used as indicators of environmental bioavailability. Whatever chemical methods are employed, none are designed per se to address the diversity of responses observed among different plant species or cultivars which can be attributed to a) the uptake behaviour of plants (i.e. sensitive, tolerant, accumulator, or hyper-accumulator of trace elements) and/or b) the ability of plants to alter the biological, physical and physical-chemical properties of their “bio-influenced zone” at the soil-root interface, i.e. the so-called rhizosphere. It could alternatively, be suggested to apply chemical methods directly to the rhizosphere but the sampling of the rhizosphere is definitely too tedious to be applied routinely.

For biological methods, four standardized biotests account for rhizosphere processes as they are based on soil-grown plants (ISO 11269-1, ISO 11269-2, ISO 17126, and ISO 22030). However, these were only designed to predict trace element phytotoxicity, i.e. the toxicological bioavailability. In these biotests, roots grow directly in the soil, therefore requiring a tedious washing procedure to reliably measure trace elements accumulated in the roots. Indeed, the amount of trace elements accumulated in shoots of non-accumulator plant species is not sufficiently sensitive to be used for assessing the environmental bioavailability of trace elements compared to the amount accumulated in the whole plant, roots included. Thus, there is still a need to develop biological methods accounting for rhizosphere processes and enabling to include the root compartment in order to properly estimate the environmental bioavailability of trace elements to plants.

Consequently, the present International Standard introduces a biotest based on the growing of roots in contact with the soil but without penetrating it. Although this experimental design is partly artificial, it enables a fair comparison of the bioavailability of trace elements between tested soils. In addition, the end point measured can be more directly related to the measurement of the environmental availability than any end point based on the measurement of toxicity.

Soil quality — Plant-based test to assess the environmental bioavailability of trace elements to plants

1 Scope

This International Standard specifies the plant-based test, hereafter called the biotest. It enables estimation of the environmental bioavailability of trace elements to plants either basically as the concentration in shoots and roots or in a more integrative way as the net uptake flux in plants. The biotest procedure includes two successive steps: (i) a pre-growth of plants in hydroponics and (ii) a growth of plants in contact with soil samples. The concentration in shoots and roots as well as the net uptake flux of trace elements in plants are determined at the end of the second step of the biotest procedure.

This biotest is applicable to the assessment of environmental bioavailability of trace elements to plants, more particularly to agricultural plants, in soils or soil materials under oxic conditions, considering that

- three plant species (cabbage, *Brassica oleracea*; tall fescue, *Festuca arundinacea*; tomato, *Lycopersicon esculentum*; 7.1) are suggested in the standardized biotest procedure, but additional target-plant species can also be used (see 7.1, Annex A), and
- the standardized biotest procedure is validated for a range of trace elements including arsenic (As), cadmium (Cd), chromium (Cr), cobalt (Co), copper (Cu), lead (Pb), nickel (Ni), and zinc (Zn), but additional trace elements can also be accounted for (see Annex A).

The biotest can be applied to soils and soil materials, including soils amended before or after field sampling with composts, sludges, wastewaters, and other (waste) materials.

NOTE 1 This biotest is not designed to assess the environmental bioavailability of trace elements that are prone to volatilisation or resulting from uptake occurring in plant leaves following, e.g. atmospheric fallout.

NOTE 2 This biotest is not designed to assess the environmental bioavailability to plants of organic contaminants. A similar experimental procedure could be used but the physical separation between plant roots and soil using a polyamide mesh needs to be adapted to avoid organic contaminant sorption on the mesh.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 3696, *Water for analytical laboratory use — Specification and test methods*

ISO 10390, *Soil quality — Determination of pH*

ISO 10694, *Soil quality — Determination of organic and total carbon after dry combustion (elementary analysis)*

ISO 11269-2, *Soil quality — Determination of the effects of pollutants on soil flora — Part 2: Effects of contaminated soil on the emergence and early growth of higher plants*

ISO 11277, *Soil quality — Determination of particle size distribution in mineral soil material — Method by sieving and sedimentation*

ISO 11465, *Soil quality — Determination of dry matter and water content on a mass basis — Gravimetric method*

This is a free preview. Purchase the entire publication at the link below:

[Product Page](#)

-
- [Looking for additional Standards? Visit Intertek Inform Infostore](#)
 - [Learn about LexConnect, All Jurisdictions, Standards referenced in Australian legislation](#)
-