



**NSAI**  
Standards

Irish Standard  
I.S. EN 16339:2013

# Ambient air - Method for the determination of the concentration of nitrogen dioxide by diffusive sampling

## I.S. EN 16339:2013

*Incorporating amendments/corrigenda/National Annexes issued since publication:*

The National Standards Authority of Ireland (NSAI) produces the following categories of formal documents:

I.S. xxx: Irish Standard – national specification based on the consensus of an expert panel and subject to public consultation.

S.R. xxx: Standard Recommendation - recommendation based on the consensus of an expert panel and subject to public consultation.

SWIFT xxx: A rapidly developed recommendatory document based on the consensus of the participants of an NSAI workshop.

*This document replaces:*

*This document is based on:*  
EN 16339:2013

*Published:*  
9 August, 2013

This document was published under the authority of the NSAI and comes into effect on:  
9 August, 2013

**ICS number:**

13.040.20

**NSAI**  
1 Swift Square,  
Northwood, Santry  
Dublin 9

T +353 1 807 3800  
F +353 1 807 3838  
E standards@nsai.ie  
W NSAI.ie

**Sales:**  
T +353 1 857 6730  
F +353 1 857 6729  
W standards.ie

Údarás um Chaighdeáin Náisiúnta na hÉireann

ICS 13.040.20

English Version

## Ambient air - Method for the determination of the concentration of nitrogen dioxide by diffusive sampling

Air ambiant - Méthode pour la détermination de la concentration du dioxyde d'azote au moyen d'échantillonneurs par diffusion

Außenluft - Bestimmung der Konzentration von Stickstoffdioxid mittels Passivsammler

This European Standard was approved by CEN on 15 June 2013.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION  
COMITÉ EUROPÉEN DE NORMALISATION  
EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: Avenue Marnix 17, B-1000 Brussels

---

<b>Contents</b>		Page
<b>Foreword</b> .....		<b>3</b>
<b>Introduction</b> .....		<b>4</b>
<b>1</b>	<b>Scope</b> .....	<b>5</b>
<b>2</b>	<b>Normative references</b> .....	<b>5</b>
<b>3</b>	<b>Terms and definitions</b> .....	<b>5</b>
<b>4</b>	<b>Principle of the method</b> .....	<b>7</b>
<b>5</b>	<b>Materials</b> .....	<b>8</b>
<b>6</b>	<b>Sampling</b> .....	<b>12</b>
<b>7</b>	<b>Analytical procedure</b> .....	<b>13</b>
<b>8</b>	<b>Calculation of the concentration of nitrogen dioxide</b> .....	<b>16</b>
<b>9</b>	<b>Quality control/quality assurance</b> .....	<b>17</b>
<b>10</b>	<b>Report</b> .....	<b>18</b>
<b>11</b>	<b>Performance requirements and measurement uncertainty</b> .....	<b>18</b>
<b>Annex A (normative) Description of samplers</b> .....		<b>21</b>
<b>Annex B (informative) Other samplers</b> .....		<b>26</b>
<b>Annex C (informative) Estimation of the uptake rate of the samplers</b> .....		<b>34</b>
<b>Annex D (informative) Measurement uncertainty</b> .....		<b>39</b>
<b>Bibliography</b> .....		<b>47</b>

## **Foreword**

This document (EN 16339:2013) has been prepared by Technical Committee CEN/TC 264 "Air quality", the secretariat of which is held by DIN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by January 2014, and conflicting national standards shall be withdrawn at the latest by January 2014.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

According to the CEN/CENELEC Internal Regulations, the national standards organisations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

## Introduction

Experience gained across the European Union (EU) in implementing EU ambient air quality legislation [1] has shown that, generally, for nitrogen dioxide (NO<sub>2</sub>), meeting the annual average limit value of 40 µg/m<sup>3</sup> is more problematic than meeting the 1-h limit value of 200 µg/m<sup>3</sup> [2].

EU Directive 2008/50/EC [1] stipulates that European Union Member States shall apply the reference measurement methods and criteria specified in the Directive. For NO<sub>2</sub> monitoring in ambient air, the reference method being that described in EN 14211:2012 [3]. However, a Member State may use any other method that provides results equivalent to that of the reference method, to be demonstrated in accordance with the Guide for the demonstration of equivalence of ambient air monitoring methods [4]. The GDE devotes specific paragraphs to methods based on diffusive sampling.

For the measurement of longer-term average concentrations of nitrogen dioxide for comparison with the annual average limit value diffusive sampling is an attractive alternative to fixed monitoring using the reference methodology described in EN 14211 because of

- small size of diffusive samplers;
- no requirement for electric power;
- potential for covering areas with a high spatial density;
- cost effectiveness.

Consequently, diffusive samplers can partially substitute and supplement fixed monitoring as an instrument for the assessment of air quality, provided that they fulfil the specific Data Quality Objectives given in [1].

At the time of publication of this standard, no full demonstration of equivalence according to [4] has been performed. However, some studies have compared NO<sub>2</sub> annual average concentrations measured by chemiluminescence and by diffusive samplers [5], [6], [7] and [8]. These have shown the potential of diffusive sampling to meet the data quality objective of 15 % expanded uncertainty for fixed measurements [1].

The methodology described in this standard can be applied to obtain air quality information with a relatively high spatial density that can be used to complement the appropriate siting of fixed monitoring stations, or in the validation of dispersion models.

Further, the methodology described can be used for simultaneously measuring sulphur dioxide (SO<sub>2</sub>) when using ion chromatography as the method of analysis. The analytical method is described in [9], [10] and [11].

This standard has been prepared based on the findings of reviews of implemented diffusive samplers in the European Union [12].

The methodology described in this standard may also be used to determine NO<sub>2</sub> in indoor air. Appropriate strategies for NO<sub>2</sub> measurement in indoor air are described in EN ISO 16000-15.

## 1 Scope

This European Standard specifies a method for the sampling and analysis of NO<sub>2</sub> in ambient air using diffusive sampling followed by extraction and analysis by colorimetry or ion chromatography (IC). It can be used for the NO<sub>2</sub> measurement in a concentration range of approximately 3 µg/m<sup>3</sup> to 130 µg/m<sup>3</sup>. A sample is typically collected for a period of 1 to 4 weeks [13], with exposure periods depending on the design of the samplers and the concentration levels of NO<sub>2</sub>.

Several sorbents can be used for trapping NO<sub>2</sub> in ambient air using a diffusive sampler. This standard specifies the application of triethanolamine as the reagent.

Nitrous acid and peroxyacetyl nitrate are the major chemical interferences of sorption by triethanolamine. However, in ambient air monitoring over long sampling times, both contaminants are generally present at low concentrations relative to NO<sub>2</sub>. Moreover, these species can also interfere with the measurement of NO<sub>2</sub> when applying the EU reference method for NO<sub>2</sub> monitoring based on chemiluminescence (see [2]).

This standard describes the application of a tube-type sampler with either a cylindrical or a slightly conical tube. Its typical uptake rate is about 1 cm<sup>3</sup>/min. Only for this sampler type sufficient evidence of validation has been found in a literature survey [12].

The relative expanded uncertainty of NO<sub>2</sub> measurements performed using these tube-type diffusive samplers can potentially be lower than 25 % for individual measurements. When aggregating results to form annual average values, the relative expanded uncertainty can be further reduced to levels below 15 % due to the reduction of random effects on uncertainty [6].

## 2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN ISO/IEC 17025, *General requirements for the competence of testing and calibration laboratories (ISO/IEC 17025)*

## 3 Terms and definitions

For the purpose of this document, the following terms and definitions apply.

### 3.1

#### **certified reference material**

reference material [3.8], characterized by a metrologically valid procedure for one or more specified properties, accompanied by a certificate that provides the value of the specified property, its associated uncertainty, and a statement of metrological traceability

[SOURCE: ISO Guide 35:2006]

### 3.2

#### **combined standard uncertainty**

standard measurement uncertainty [3.10] that is obtained using the individual standard measurement uncertainties associated with the input quantities in a measurement model

[SOURCE: JGCM 200:2012]

### 3.3

#### **desorption efficiency**

ratio of the mass of analyte desorbed from a sampling device to that applied

This is a free preview. Purchase the entire publication at the link below:

[Product Page](#)

- 
- [Looking for additional Standards? Visit Intertek Inform Infostore](#)
  - [Learn about LexConnect, All Jurisdictions, Standards referenced in Australian legislation](#)
-