



NSAI
Standards

Irish Standard
I.S. EN 16618:2015

Food analysis - Determination of acrylamide in food by liquid chromatography tandem mass spectrometry (LC-ESI-MS/MS)

I.S. EN 16618:2015

Incorporating amendments/corrigenda/National Annexes issued since publication:

The National Standards Authority of Ireland (NSAI) produces the following categories of formal documents:

I.S. xxx: Irish Standard — national specification based on the consensus of an expert panel and subject to public consultation.

S.R. xxx: Standard Recommendation — recommendation based on the consensus of an expert panel and subject to public consultation.

SWiFT xxx: A rapidly developed recommendatory document based on the consensus of the participants of an NSAI workshop.

This document replaces/revises/consolidates the NSAI adoption of the document(s) indicated on the CEN/CENELEC cover/Foreword and the following National document(s):

NOTE: The date of any NSAI previous adoption may not match the date of its original CEN/CENELEC document.

This document is based on:

EN 16618:2015

Published:

2015-04-22

This document was published under the authority of the NSAI and comes into effect on:

2015-05-09

ICS number:

67.050

67.060

67.080.20

67.140.20

NOTE: If blank see CEN/CENELEC cover page

NSAI
1 Swift Square,
Northwood, Santry
Dublin 9

T +353 1 807 3800
F +353 1 807 3838
E standards@nsai.ie
W NSAI.ie

Sales:
T +353 1 857 6730
F +353 1 857 6729
W standards.ie

Údarás um Chaighdeáin Náisiúnta na hÉireann

EUROPEAN STANDARD

EN 16618

NORME EUROPÉENNE

EUROPÄISCHE NORM

April 2015

ICS 67.050; 67.060; 67.080.20; 67.140.20

English Version

Food analysis - Determination of acrylamide in food by liquid chromatography tandem mass spectrometry (LC-ESI-MS/MS)

Analyse des produits alimentaires - Dosage de l'acrylamide dans les produits alimentaires par chromatographie en phase liquide couplée à la spectrométrie de masse en tandem (CL-ESI-SM-SM)

Lebensmittelanalytik - Bestimmung von Acrylamid in Lebensmitteln mit Flüssigchromatographie und Tandem-Massenspektrometrie (LC-ESI-MS/MS)

This European Standard was approved by CEN on 7 February 2015.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

Contents		Page
Foreword		3
1	Scope	4
2	Normative references	4
3	Principle	4
4	Reagents	4
5	Apparatus	7
6	Sample preparation	8
7	Measurement	10
8	Determination of concentrations	13
9	Precision	15
10	Test report	17
Annex A (informative) Typical chromatograms		18
Annex B (informative) Precision data		20
Bibliography		24

Foreword

This document (EN 16618:2015) has been prepared by Technical Committee CEN/TC 275 “Food analysis - Horizontal methods”, the secretariat of which is held by DIN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by October 2015 and conflicting national standards shall be withdrawn at the latest by October 2015.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

This document has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association.

WARNING — The use of this document can involve hazardous materials, operations and equipment. This document does not purport to address all the safety problems associated with its use. It is the responsibility of the user of this document to establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

EN 16618:2015 (E)

1 Scope

This European Standard specifies a method for the determination of acrylamide in bakery ware such as bread, toasted bread, crisp bread, butter cookies, and biscuits, as well as potato products such as potato chips, potato crisps, and potato pan cake and roasted coffee, by liquid chromatography in combination with electrospray ionization and tandem mass spectrometry (LC-ESI-MS/MS). This method has been validated in an interlaboratory study via the analysis of both naturally contaminated and spiked samples, ranging from 14,3 µg/kg to 9 083 µg/kg. It was developed at the Swedish National Food Administration and validated in a study organized by the Directorate General Joint Research Centre (DG JRC), Swedish National Food Administration and the Nordic Committee on Food Analysis (NMKL), see [1] and [2].

The limit of quantification (LOQ) depends on the type of instrument used and on the actual performance of the instrument. The majority of the laboratories participating in the validation study were able to determine acrylamide in a butter cookie sample at a level of 14,3 µg/kg. Thus, the validation by interlaboratory study showed that LOQ can be expected to be in the range between below 15 µg/kg and 30 µg/kg.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN ISO 1042:1999, *Laboratory glassware - One-mark volumetric flasks (ISO 1042:1998)*

EN ISO 3696:1995, *Water for analytical laboratory use - Specification and test methods (ISO 3696:1987)*

3 Principle

Acrylamide is extracted with water and isotopic labelled acrylamide is added. The extract is centrifuged and the supernatant is cleaned up with two solid phase extraction (SPE) columns. The first SPE column contains silica based C18 groups as well as anion and cation exchangers, and since acrylamide is not retained by the column, the extract is just passed and collected. The reason for using this column is to retain as many matrix components as possible (non-polar compounds as well as anions and cations) without retaining acrylamide, i.e. this first SPE column is used as a chemical filter.

The second SPE column contains a polymer based phase with a relatively high capacity to bind acrylamide. The extract is loaded onto the column, the column is washed with water and finally eluted with a mixture of 60 parts per volume of methanol and 40 parts per volume of water. The purpose of this step, apart from further cleaning of the extract, is to concentrate the extract and to obtain low limits of quantification.

After evaporation of the methanol, the extract is analysed by LC-MS/MS. For this purpose an HPLC column with graphitized carbon as stationary phase is used, since the retention factor (k) is relatively high ($k = 4$ when no organic solvent is added in the mobile phase) compared to other commercially available columns.

4 Reagents

Use only reagents of recognized analytical grade and water complying with grade 1 of EN ISO 3696:1995, unless otherwise specified. Solvents shall be of quality for HPLC analysis.

4.1 Acrylamide (CAS 79-06-1), purity not less than 99,9 % mass fraction.

This is a free preview. Purchase the entire publication at the link below:

[Product Page](#)

-
- [Looking for additional Standards? Visit Intertek Inform Infostore](#)
 - [Learn about LexConnect, All Jurisdictions, Standards referenced in Australian legislation](#)
-